

স্কাবিড

পারমিথ্রিন বিপি ৫% ডব্লিউ/ডব্লিউ

বর্ণনাঃ

~vveW n#Q cvBwi# ~vBwM RvRK| Bnv n#Q mv'v my#Kvgjxq cvwb l wgwKÖbxq
wju hmv evwn'K e'env#ii Rb']

উপাদানঃ

প্রতি ১০০ গ্রাম ক্রিমে আছেঃ

cviwgv_#b wewc 5 MÖvg

নির্দেশনাঃ

~vwem Ges μ'~ve jvBm Bb#dm#Ukb Gi wPwKrmvq wbf'©wkZ|

মাঝা ও প্রয়োগবিধিঃ

evwn'Kfv#e cÖ#qvM Ki#Z n#e|

wbgœwjlwZ tWv#RR wmwWDj Abzmv#i e'envi Ki#Z n#e |

c~Y@eq r l 12 erm#ii E#aY@ t 30 MÖvg ch@š|

wki#i t#i#i 6-12 ermi t15 MÖvg ch@š|

1- 5 ermi ch@š| t 7 MÖvg ch@š|

2 gvm- 1 ermi t 4 MÖvg ch@š|

পদ্ধতি ও প্রয়োগঃ

ZjK cwi'vi Kiæb Ges cÖ#qv#Mi c~#e@ fvj fv#e i'®. Ges kxZj Kiæb| Zvi c#i ~vveW
wju Øviv gv_v e'ZxZ mg' l kix#i jvMvb| 8 - 12 NÈv ci fvj fv#e gv_v e'ZxZ
cÖ#qvMK...Z ~vb ay#q fdjyb| mvaviYZ ~vwem Bb#dKkb GK evi e'env#iB
AcmwviZ nq| wKš' ÖcÖ_g cÖ#qv#Mi mvZ w'b ci c~bivq e'envi Gi cÖ#qvRb n#Z
cv#i| 12 eQ#ii w#gœ wki#i Ges eq'ç'i Rb' mZK@Zvi mv#_ Kv#iv mycvi wfk#b
e'envi Kiv DwPZ|

পার্শ্ব প্রতিক্রিয়াঃ

Zj#Ki weiaç fve thgbt cÖ'vn n#Z cv#i| Zj#Ki PzjKvwbnm BivB#ngv, BwWgv, i'vk
t'Lv w'#Z cv#i |

প্রতিনির্দেশনাঃ

cviwgv_#bi cÖwZ th mKj tivMx#i m#e'bkxjZv i#q#Q Zv#i t#i#i cÖwZ
wb#i'©kbv i#q#Q |

পূর্ব সতর্কতাঃ

t'PvL I gy#L ~vveW thb bv jv#M t#m w'#K mZK@Zv Aej#^b Ki#Z n#e|

সতর্কতাঃ

th mKj e'w³ tewk tewk cviwgv_#b e'envi K#i Zv#i nv#Zi PzjKvwb Kgv#bvi Rb'
t'Movfm e'envi Kiv DwPZ|

গর্ভাবস্থায় ও স্তন্যদানকালে ব্যবহারঃ

Mf@ve'vq Wv'v#ii civgK@ Abyhvqx e'envi Ki#Z n#e| th#nZz Mf@ve'vq Gi we#kl
t'Kvb cixyvi cÖgvY t#b| 'y» vbKv#j l gvZ... y#x cviwgv_#bi Dcw 'wZ m#u#K@
Rvbv hvqwb, ZvB GwU 'y» vbKvix gv#q#i t#i#i e'envi Kiv DwPZ bq|

জন্ম গ্ৰহণের সাথে আন্তঃক্রিয়াঃ

~vveW wgv_#Zgb Rvbv t#b| KwU@#Kv#++vi#qW t#m#b Ki#j cviwgv_#b e'envi Kiv
DwPZ |
Chemist
কলেজ রো, বরিশাল, বাংলাদেশ।

SCABID

Permethrin BP 5% w/w

DESCRIPTION:

SCABID is a recently introduced pyrethroids derivative. It is a white, smooth and water miscible cream intended for external application.

COMPOSITION:

Each 100 gm cream contains:

Permethrin BP 5 gm

INDICATIONS:

Treatment of scabies and crab lice infestations.

DOSAGE AND ADMINISTRATION: External application.

The following dosage schedule is given for a single application. Adult and children over

12 years Children: : up to 30 gm

6-12 years : up to 15 gm

1-5 years : up to 7 gm

2 month 1 year : up to 4 gm

PROCEDURE FOR APPLICATION:

Clean the skin, dry and cool before applying SCABID. Rub gently SCABID over the whole body excluding head. Then wash thoroughly after 8-12 hours. Do not wash with soap during treatment. Normally the scabies infection is cleared in a single application. But a reapplication may be necessary after seven days of the first application. Children under 12 years and old patients should be carefully supervised by adults.

SIDE-EFFECTS:

Skin discomfort e.g. burning or stinging usually occurs in patients with severe scabies. Other side-effects are skin irritation including erythema, oedema, eczema, rash and pruritus.

CONTRAINDICATIONS:

SCABID is contraindicated in patients known to be hypersensitive to permethrin, other pyrethroids or pyrethrins.

PRECAUTION:

Avoid contact of SCABID with the eyes and mouth.

WARNING:

Persons frequently coming in contact with permethrin should wear gloves to avoid possible irritation to the hands.

USE IN PREGNANCY & LACTATION:

In the absence of specific studies in pregnant women, its use in pregnancy should only follow medical advice. It is not known whether permethrin is excreted in human milk so it should not be used in nursing mothers.

DRUG INTERACTION:

There is no known drug interaction with permethrin. However treatment with corticosteroids in eczematous conditions should be stopped during application of permethrin.

STORAGE:

Store at temperature within 30°C, protected from light & moisture. Keep out of the reach of children.

PACKAGE QUANTITIES:

Tubes containing 15 gm or 30 gm of cream.

Manufactured by:

Chemist Laboratories Ltd.

College Row, Barishal, Bangladesh.