


 <h2 style="text-align: center;">এনজিট্যান</h2> <p style="text-align: center;">tjvmviUvb cUvwkqvg U`ve#jU</p>	 <h2 style="text-align: center;">Angitan</h2> <p style="text-align: center;">Losartan Potassium BP 50 mg</p>
<p>উপাদান cÖwZwU U`ve#j#U i#q#Q tjvmviUvb cUvwkqvg wewc 50 wg.MÖv. </p> <p>বিবরণ এনজিট্যান (tjvmviUvb cUvwkqvg) n#`Q GKwU bZzb cÖR#b#i Gw>UnvBcvi#UwYf W^vM-Gi cÖ_g W^vM, hv GwÄI#Ubwmb II wi#m#i (UvBc GwU1) Gi G>UvMwb#i GwÄI#Ubwmb II GKwU AZ`í Kvh©Kix f`v#mvKYwU^KUi hv tiwbb GwÄI#Ubwmb wm#:# Gi cÖv_wgK f`v#mvGKwUf ni#gvb Bnv nvBcvi#Ubk c`v#_vwdwRljwRi GKwU ,iæZ;c~Y© Dcv`vb tjvmviUvb Zvi cÖavb Kvh©Ki tgUv#evjvBU, GwÄI#Ubwmb II-Gi Øviv m,ó i#bvjx ms#KvPb I Gj#Wv#:#ib wbtmib cÖwZ#iva K#i tjvmviUvb mywbw`©ófv#e GwÄI#Ubwmb II-Gi mv#_ GwU1 wi#m#i Gi evBwÜs cÖwZnZ K#i KvR K#i এনজিট্যান GLb D`P i#Pv#ci wPwKrmvq cÖv_wgK Jla wn#m#e `^xK...Z </p> <p>নির্দেশনা এনজিট্যান mKj cÖKvi D`P i#Pvc Ges Kb#RmwUf nvU© tdBwJli - G e`eüZ nq Bnv GKKfv#e A_ev Ab`vb` D`P i#Pvc wbq#z#bKvix lly#ai mv#_ e`envi Kiv th#Z cv#i GwmB BbwnweUi tme#bi ci m,ó i` Kvwki d#j llya tmeb eÜ nlqvi t#j#i এনজিট্যান GKwU Kvh©Kix weKij সেবন মাত্রা ও বিধি এনজিট্যান -Gi `^vfvweKgv#v `wbK 1wU U`ve#jU 25 wg.MÖv. (A#a©K U`ve#jU) e`envi Kiv nq tm mg`I tivMxi t#i hv#`i B>U^vfmKzjvi fj`g wWwçøkb -Gi m#çvebv i#q#Q (thgb-WvBBD#iwUK MÖnYKvix) Ges hv#`i hK...#Zi Amg Kvh©KvwiZvi BwZnm Av#Q tjvmviUvb `wbK 25-50 wg.MÖv. GK t_#K `yÖev#i t`qv th#Z cv#i `wbK GKevi tme#b hw` Gw>UnvBcvi#Ubwmbf B#d± Ach©vß g#b nq tm#j#i `wbK tgUv gvlv Awfboe ti#L wKsev wKQzUv evwo#q cÖwZw`b `yÖevi e`envi Kiv hv#e hw` D`P i#Pvc tKejgvl tjvmviUvb e`env#ii d#j wbq#w#Z bv nq Z#e wb#ægv#vq WvBBD#iwUK mn#hv#M e`envi#hvM` Bnv Lvev#ii mv#_ A_ev Avjv`vfv#el Lvlqv th#Z cv#i </p> <p>পার্শ্ব-প্রতিক্রিয়া tjvmviUvb cUvwkqvg Gi cvk ©-cÖwZwµqv LyeB Kg hv çøvw#ev -Gi m#½ Zzjbxq çøvw#ev -Gi PvB#Z 1% telx tivMxi t#j#i N#U Ggb cvk ©-cÖwZwµqv g#a` Avcv#i timwç#iUwi U^v± Bb#dK&kb (7.9% ebvg 6.9%), gv_v wSgwSg Kiv (3.5% ebvg 2.1%) Ges cv#q e`_v (1.0% ebvg 0.0%) D#jØL#hvM` </p> <p>সতর্কতা tiwbb-GwÄI#Ubwmb wm#:#-Gi gva`fg KvR K#i Ggb Jla Gi cÖwZ hviv `úk©KvZi Zv#`i t#j#i tjvmviUvb cUvwkqvg mZK©Zvi m#½ e`envi Ki#Z n#e we#kl mZK©Zv Aeja^b Ki#Z n#e e,° Ges hK...#Zi Amg Kvh©KvwiZvi t#j#i wki#`i t#j#i tjvmviUvb cUvwkqvg Gi wbivc#v Ges Kvh©KvwiZv cÖwZwóZ nqwb </p> <p>প্রতিনির্দেশনা thmg`I tivMx tjvmviUvb cUvwkqvg ev GB U`ve#j#Ui tKvb Dcv`v#bi cÖwZ AwZ ms#e`bkxj Zv#`i t#j#i Gi e`envi wbwLx </p> <p>গর্ভাবস্থায় ও স্তন্যদানকালে ব্যবহার Mf©aviY wbwðZ nlqv gv#B `æZ tjvmviUvb cUvwkqvg tmeb eÜ Ki#Z n#e `lb`vbKv#j tjvmviUvb cUvwkqvg e`envi Kiv hv#e bv KviY gvZ...`y» Gi mwnZ GwU wbtm,Z nq wKbv Zvi tKvb Z` t#b </p> <p>cÖ`ZKviK :  কেমিস্ট ল্যাবরেটরীজ লিমিটেড কলেজ রো, বরিশাল, বাংলাদেশ।</p>	<p>Composition Each tablet contains Losartan Potassium BP 50 mg.</p> <p>Description Angitan (Losartan Potassium), the first of a new class of antihypertensives, is an angiotensin II receptor (type AT1) antagonist. Angiotensin II is a potent vasoconstrictor, the primary vasoactive hormone of the renin-angiotensin system and an important component in the pathophysiology of hypertension. Losartan and its principal active metabolite block the vasoconstriction and aldosterone secreting effects of angiotensin II by selectively blocking the binding of angiotensin II to the AT1 receptor found in many tissues. Angitan is now regarded as the first-line therapy option for treating high blood pressure.</p> <p>Indication Angitan is indicated for the treatment of all grades of hypertension and congestive heart failure. It may be used alone or in combination with other antihypertensive agents. It is an effective alternative for patients who have to discontinue an ACE inhibitor because of persistent dry cough.</p> <p>Dosage & Administration The usual starting dose is 1 tablet once daily. In patients with possible depletion of intravascular volume or patients with a history of hepatic impairment, starting dose is 25 mg (half tablet) once daily. Losartan can be administered once or twice daily with total daily doses ranging from 25 mg to 100 mg. No initial dosage adjustment is necessary for elderly or renal impairment patients. If blood pressure is not controlled by Losartan, a low dose of a diuretic (Hydrochlorothiazide) may be added. Angitan may be administered with or without food.</p> <p>Side Effect Overall incidence of adverse effects of Losartan potassium is comparable to placebo in clinical studies. The most common adverse events occurring with Losartan potassium at a rate of >1% above placebo were upper respiratory infection (7.9% vs 6.9%), dizziness (3.5% vs 2.1%) and leg pain (1.0% vs 0.0%).</p> <p>Precautions Losartan potassium should be used with caution in patients with known hypersensitivity to the drugs that act through renin-angiotensin system. Special precaution should be taken when it is administered to the patients with renal and hepatic impairment. Safety and effectiveness of Losartan potassium in pediatric patients have not been established.</p> <p>Contraindications Losartan potassium is contraindicated in patients who are hypersensitive to the active ingredient or any component of the drug.</p> <p>Use in Pregnancy & Lactation Losartan potassium must be discontinued as soon as possible when pregnancy is detected. It should not be prescribed during lactation as there is no information in humans on the passage of Losartan (Angitan) into breast milk.</p> <p>Drug Interaction No drug interactions of clinical significance have been identified. Drugs which have been studied in clinical pharmacokinetic trials include - (1) Hydrochlorothiazide, (2) Digoxin, (3) Warfarin, (4) Cimetidine, (5) Ketoconazole and (6) Phenobarbital.</p> <p>Storage Store at temperature within 30°C, protected from light & moisture.</p> <p>Commercial Packaging Angitan tablet: Each box containing 3 Alu-Alu strips of 10 tablets.</p> <p>Manufactured by :  Chemist Laboratories Ltd. College Row, Barishal, Bangladesh.</p>