

উইনকফ

Gg#e#v-j nvB#W#v#K-vivBW wewc

উপাদান

উইনকফ wmicv: cÖwZ 5 wg.wj. wmicv# Av#Q Gg#e#v-j nvB#W#v#K-vivBW wewc 15 wg.MÖv.]

ফার্মাকোলজি

Gg#e#v-j te#vg#nw-b-Gi mwµq tgUv#evjvBU Ges te#vg#nw-b-Gi tP#q Gi Kvh@KwviZv tekx| nvB#W#v#Kb c#wµqvi gva`#g t#k#Sv#K k#vmbvjxi g#a` Zij K#i Ges k#vMkó Kgvq| Gg#e#v-j Gjwfljvi tKv#li mvi#dK#U-U-Gi ,iæZ;c~Y© Dcv`vb dm#dvwjwvW ^Zix Z;ivwš^Z K#i, d#j GjwfljvB-Gi wfZ#ii Pvc K#g| GUv e#w¼qvj nvBcviGw±wfwU Kgvq| Gg#e#v-j mvB#UvKvBb Ges GivwKWwbK GwmW tgUv#evjvBU ^Zix#Z evav w`#q cÖ`vn#ivax f,wgKv iv#L| COPD tivMx#i k#vmbvjxi cÖk`Izv eRvq ivL#Z mvnvh` K#i|

নির্দেশনা

- t#k#Svhy³ k#vMz#šzi GwKDU I µwYK cÖ`vn thgb GwKDU I µwYK e#¼vBwUm
- t#k#Svhy³ ivB#bv#dwiv#Äqvj U#v±-Gi cÖ`vn (j`vwib&RvBwUm, d`vwib&RvBwUm mvBbmvBwUm Ges ivBbvBwUm)
- t#k#Svhy³ GRgvwUK e#¼vBwUm, e#w¼qvj G`vRgv
- e#w¼qvjK#Uwmm
- µwYK wbD#gwbqv|

সেবনবিধি ও মাত্রা

WbK tmeb gvUv (Lvevi ci MÖNY Kiv DĒg):

উইনকফ wmicv

2-5 eQi eqm : 2.5 wg.wj. (1/2 Pv PvgP) w`#b

2-3 evi

5-10 eQi eqm : 5 wg.wj. (1 Pv PvgP) w`#b 2-3

evi

10 eQ#ii tewk eq#mi wki l eq`c

: 10 wg.wj. (2 Pv PvgP) w`#b 2-3 evi

পার্শ্ব প্রতিক্রিয়া

wKQy wKQZ t#j#i cwicvKZ#šzi wewfbœ cvk|© cÖwZwµq v thgb cvK`jxi cÖ`vn I fvi fvi fve t`Lv w`#Z cv#i| K`vwPr GjvwR©, thgb- PzjKvwb, GbwRlwbD#ivwUK B#Wgv n#Z cv#i|

প্রতিনির্দেশনা

Gg#e#v-j ev te#vg#nw-b Gi AwZms#e`bkxjZv_vK#j|

সাবধানতা

M`vw±#K I wWl#Wbvj Avjmv#ii tivMx#i Ges wLuPzbx i tivMx#i mveavbZvi mv#_ Gg#e#v-j t`qv DwPZ| hv#i hK...Z I e.,#i Kvh#ygvZvi Ach#vβZv i#q#Q Zv#i t#j#i mveavbZv Aej#Ab Kiv DwPZ|

অন্য ওষুধের সাথে প্রতিক্রিয়া

Gw>Uuzwmf (thgb tKv#WbB) Gi mv#_ MÖNY Kiv DwPZ bq, t#j#i t#k#Sv hv Gg#e#v-j Øviv Zij n#q#Q Zv t#i n#q th#Z cv#i bv|

গর্ভাবস্থায় ও স্তন্যদানকালে ব্যবহার



কেমিস্ট ল্যাবরেটরীজ লিমিটেড
কলেজ রো, বরিশাল, বাংলাদেশ।

Winkof

Ambroxol Hydrochloride BP

COMPOSITION

Winkof Syrup (100 ml) : Each 5 ml of Syrup contains 15 mg Ambroxol Hydrochloride BP.

PHARMACOLOGY

Winkof is the active metabolite of bromhexine and it has been proven that this metabolite possesses a greater bronchosecretolytic effect than bromhexine. It improves sputum rheology by hydrating mechanism leading to liquefaction of mucus in the lumen of respiratory tract, thus facilitating expectoration of mucus and reducing dyspnea. It stimulates production of phospholipids of surfactant by alveolar cells, thus contributing to the lowering of superficial tension in the alveoli. It also reduces bronchial hyperactivity. Winkof has anti-inflammatory properties owing to the inhibitory effect on the production of cellular cytokines and arachidonic acid metabolites. In patients with COPD traditionally improves airway potency.

INDICATIONS

Productive cough, Acute and chronic inflammatory disorders of upper and lower respiratory tracts associated with viscid mucus including acute and chronic bronchitis, Inflammatory disease of rhinopharyngeal tract (laryngitis, pharyngitis, sinusitis and rhinitis) associated with viscid mucus, Asthmatic bronchitis, bronchial asthma with thick expectoration, Bronchiectasis, Chronic pneumonia.

DOSAGE AND ADMINISTRATION

Average daily dose (preferably after meal):

Winkof syrup:

2-5 years old : 2.5 ml (1/2 teaspoonful), 2-3 times a day

5-10 years old : 5 ml (1 teaspoonful), 2-3 times a day

10 years old and adults: 10 ml (2 teaspoonful) 2-3 times a day

SIDE EFFECTS

Gastrointestinal side effects like epigastric pain, stomach overfill feeling may occur occasionally. Rarely allergic responses such as eruption, urticaria or angioneurotic edema has been reported.

CONTRAINDICATIONS

Contraindicated in known hypersensitivity to Winkof or Bromhexine.

PRECAUTIONS

Winkof should be given cautiously to patients with gastric and duodenal ulceration or convulsive disorders. Patients with hepatic and renal insufficiency should take it with caution.

DRUG INTERACTION

Winkof should not be taken simultaneously with antitussives (e.g. Codeine) because phlegm, which has been liquefied by Winkof might be expectorated.

USE IN PREGNANCY AND LACTATION

Teratogenic and foetal toxicity studies have shown no harmful effect of Winkof however, it is advised not to use it in pregnancy, especially during the 1st trimester. Safety during lactation has not been established yet.

STORAGE CONDITION

Protect from direct light exposure. Store in dry place at a temperature not exceeding 30°C. Keep out of the reach of children.

HOW SUPPLIED

Winkof syrup : PET Bottle contains 100ml syrup & a measuring cup.



Chemist Laboratories Ltd.

College Row, Barishal, Bangladesh.