

Tienium

Tiemonium Methylsulphate INN

Composition:

Tienium 50 mg tablet:

Each film coated tablet contains 50 mg of Tiemonium Methylsulphate INN.

Tienium 5 mg/2 ml injection:

Each 2ml ampoule contains 5 mg of Tiemonium Methylsulphate INN.

Pharmacology:

Tienium is a quaternary ammonium antimuscarinic agent with peripheral effect similar to those of atropine. Tienium strengthens calcium bonding with phospholipids and proteins thus stabilizing the cell membrane of the GI tract.

Indication and usage:

Tienium is an antispasmodic drug. It reduces muscle spasms of the intestine, biliary system, uterus & urinary bladder. It is indicated for the pain in gastrointestinal & biliary disease in the urology and gynecology such as gastroenteritis, diarrhea, dysentery, biliary colic, enterocolitis, cholecystitis, colonopathy, mild cystitis, & spasmodic dysmenorrhoea.

Dose & administration:

Route of administration: Oral. Recommended oral dose of Tienium tablet is 2-6 tablets (100-300 mg) daily in divided dose as required. Recommended injectable dose is one Tienonium Methylsulphate injection three times daily, through intravenous routeslowly or intramuscular route.

Contraindication:

Tiemonium Methylsulphate tablet should not be used in Glucoma or where acute pain of eyeball with vision disturbance was found. Also this drug should not be used in patients with the disorder of prostate or urinary bladder.

Warning and precaution:

Precautions should be exercised for treatment of patients with disorders of prostate, chronic bronchitis, coronary insufficiency, ambient hyperthermia, cardiac disorders, thyroid disorders, serious renal & hepatic insufficiency.

Side effects:

Side effect with the use of Tiemonium Methylsulphate tablet is very rare.

Use in pregnancy & lactation:

Tiemonium Methylsulphate tablet may be used in pregnancy only if it is clearly needed by the assessment of risk benefit ratio. Tiemonium Methylsulphate tablet may be used in lactating mother only if it is clearly needed by the assessment of risk benefit ratio.

Use in geriatric:

Efficacy and safety were maintained with increasing age.

Drug interaction:

Tiemonium Methylsulphate tablet should not be used with other drugs without prior consult of a registered physician to avoid possible drug interactions.

Overdose:

There is no available data regarding the overdose of Tiemonium Methylsulphate.

Storage:

Store in cool & dry place below 30°C, protect from light & moisture. Keep the medicine out of the reach of children.

Commercial Pack:

Tienium 50 mg tablet: Each box contains 5x10's tablets in blister pack.

Tienium 5 mg/2 ml injection: Each box containing 5 ampoules of 2 mg.

Manufactured by:



Chemist Laboratories Ltd.
College Row, Barishal, Bangladesh.

টাইনাম

UvB#gvwbbqvg wg_vBjmvj#dU AvBGbGb

উপাদানঃ

টাইনাম 50 wgMÖv U`ve#jUt cÖwZ wdj# +Kv#UW U`ve#j#U i#q#Q UvB#gvwbbqvg wg_vBjmvj#dU AvBGbGb 50 wgMÖv| টাইনাম 5 wgMÖv / 2 wgwj Bb#RkKbt cÖwZ 2 wgwj G`v#úy#j Av#Q UvB#gvwbbqvg wg_vBjmvj#dU AvBGbGb 5 wgMÖv| ফার্মাকোলজিঃ

টাইনাম GKwU †KvqvU@vibvix A`v#gvwbbqvg Dcv`vb hvi A`v#U#vwc#bi gZ KvH@KvwiZv i#q#Q| টাইনাম dm#dwjwcW I †cÖvUwB †gg#e#bi K`vjwmqvg eÜb kw`kvjx K#i hvi d#j cwicvKZ#š#i †Kvl AveiYx w`vZkxj nq|

নির্দেশনাঃ

টাইনাম GKwU Gw`U`úvR#gvwWK lly| Bnv cwicvKZš#i, wewjqvix wm#:#g, BD#Uivm I g`~lvjxi †cxi ms#KvPb#K wkw_j K#i| Bnv M`v#:#vG`U#vBwUm, Wvqvwiqv, wW#mw#U#v, wewjqvix KwjK, G`Uv#iv#KvjvBwUm, Kwjwm#vBwUm, †Kv#jv#bvc`v_x, nvjKv wm#vBwUm Ges `úvR#gvwWK wWm#g#bvwiqv#Z wb#`@wkZ|

মাত্রা ও সেবনবিধিঃ

†meb Kivi cxwZt gy#L †mbe †hvm`| টাইনাম U`ve#jUt 2-6 wU U`ve#jU (100-300 wgMÖv) `wbK we#`gvlvq gy#L †L#Z n#e|

টাইনাম Bb#RkKbt 1 wU K#i G`v#úy#j (5 wgMÖv) w`#b 3 evi B`U#v#gvmKzjvi A_ev B`U#v#fbvm c#_ax#i ax#i cÖ#qvM Ki#Z n#e||

প্রতিনির্দেশনাঃ

UvB#gvwbbqvg wg_vBjmvj#dU Møy#Kvgv tivMx#`i †#i#f Ges †h mKj tivMxi GwKDU AvBej †cBb I †mB mv#_`_,wói mgm`v Av#Q Zv#`i †#i#f e`envi Kiv hv#e bv| hv#`i †cÖv#:#U A_ev BDwibvix eovWv#ii A`#vfvweKZv Av#Q Zv#`i †#i#f Bnv e`envi Kiv hv#e bv|

সতর্কতাঃ

cÖ#:#Ui tivM, `xN@`vqx e#%4vBwUm, K#ivbwwi Ac~Y@Zv, cwi#ewöZ nvBcvi_vwg@qv, ü`#ivM RwbZ mgm`v, _vBi#qW RwbZ mgm`v, ,iæZi †lwZKvik Ges †ncvwUK Ac~Y@Zv tivMx#`i wPwKrmvi Rb`mZK@Zv Aeja#^b Kiv DwPZ|

পার্শ্ব-প্রতিক্রিয়াঃ

`#vfeZ LyeB Ai cwigvY cvk#`cÖwZwµqv †`Lv hvq|

গর্ভাবস্থায় ও স্তন্যদানকালে ব্যবহারঃ

ãæ#Yi †lwZ A#cvlv gv#qi DcK...Z nevi m#cvbev tewk _vK#j e`envi Kiv hv#e|

mšlv#bi †lwZ A#cvlv gv#qi DcK...Z nevi m#cvbev tewk _vK#j e`envi Kiv hv#e|

বয়স্কদের ক্ষেত্রে ব্যবহারঃ

Kvh@KvwiZv Ges wbivcEv µgea#gvb eq#mi mv#_ evRvq _v#K|

অন্য ওষুধের সাথে প্রতিক্রিয়াঃ

wPwKrm#Ki civgk@ e`ZxZ Ab` lly#ai mv#_ UvB#gvwbbqvg

cÖ`ZKvik :



কেমিস্ট ল্যাবরেটরীজ লিমিটেড
কলেজ রো, বরিশাল, বাংলাদেশ।