

পেইনেক্স

U`vefjU

Gwm#K-v#dbvk wewc

উপাদান :

cOwZ wdjY-fKv#UW U`vefjU Av#Q Gwm#K-v#dbvk wewc 100 wg.MÖv.

বর্ণনা :

t#b#b- Gwm#K-v#dbvk i#q#Q hv GKwU bb-fU#ivBwj cÖ`vn Ges e`_v bvkK G#R#U| Bnvi Kv#KvwiZv eüjvs#k t#CÖv-v#mÖvWb Drcv`#bi ms#M RwoZ mvB#K-v#Aw-f#R#bi ms#kol#Y evav cÖ`v#bi Dci wbf#ikxj| Bnv ZiæYw` MV#b cÖ`q#Rbxq t#gSwjk c`v`© MÖvBK`vbm Gi ms#kolY#Ki DrmwvwnZ K#j|

নির্দেশনা :

t#b#b- Aw-lAv_@ivBwUm, wiDgvU#qW Av_@ivBwUm Ges GbKvB#jvwRs `übwWjvBwUm RwbZ e`_v Ges cÖ`vn Dck#gi Rb`wb#`©wkZ|

মাত্রা এবং ব্যবহার বিধি :

gy#L fL#Z n#e|

eq`#`Rb` t` 1wU U`vefjU (100 wgt.MÖvt) K#i cÖWZw`b`yB evi| wki#`i Rb` t` wki#`i tiv#M Gwm#K-v#dbvk e`env#ii Rb` wK-wbK`v#j ch#v#Z`_`tb#j| ZvB wki#`i Gwm#K-v#dbvk e`envi Kiv DwPz bq| t#CÖSp#`i Rb` t` t#CÖSp tivMx#`i Rb` Gwm#K-v#dbvk#Ki t#KvB cwieZ#b b#vB| myZivs g#lv e`env#ii nvi#K cwieZ#b Kivi cÖ#qvRb we#ePbv Kiv nqw#| hK...#Zi mgm`vq` t` thme tivMx#`i hK...#Zi ywZ mwvaZ n#q#Q Zv#`i t#j#i t#b#b-i g#lv| KgvtZ n#e| cÖviva#K g#lv `wbK 1wU U`vefjU (100 wgt.MÖvt) wnmv#e e`envi Ki#Z n#e| wKWbxi mgm`vq` : wKWbxi mgv#b` ywZ mwvaZ n#q#Q tivMx#`i t#j#i Gi#f cÖgvY t#b#b i g#lv wKQzUv cwieZ#b Ki#Z n#e|

পার্শ্ব-প্রতিক্রিয়া :

t#b#b- mvaviZt mymnbxq| AwaKvsk cvk#©-cÖWZw#qv`#jv mnbxq| G me cvk#©-cÖWZw#qv`#jv n#`Q ewg ewg fve, Wvqvwiq, ARxY#Zv Ges Z`t`cU e`_v|

বিরূপ-প্রতিক্রিয়া :

thme tivMxMY Gwm#K-v#dbvk A_ev Gmwcvib A_ev Ab`vb` bb-#ö#ivBwvj cÖ`vn bvkK J#ai cÖWZ AwaK ms#e`bKvZiv t#b#b- e`envi Ki#eb bv|

গর্ভাবস্থায় ব্যবহার বিধি :

Mf#ve`vq Gwm#K-v#dbvk#i w#vive` e`envi m#u#K#© t#KvB Z`_`Rv#v b#vB| t#mRb` Mf#ve`vq GKv#l Riseix KviY Qvov t#b#b- e`envi Kiv DwPr bq|

মাতৃদুগ্ধ দানকালে ব্যবহার বিধি :

gZ...y#» Gwm#K-v#dbvk#i w#vive` m#u#K#© t#KvB Z`_`Rv#v b#vB| t#mRb` gZ...y#» `vbkv#j t#b#b- e`envi e#R#öb Ki#Z n#e` hZy` ch#l wki#`i Dc#i ma#ve` SzuuKi t#P#q gvt#`i Rx#e#i Rb` cÖ`Qb#e Dckvi AwaKZi `iæZ#-Y`© nq|

সতর্কতা :

thme tivMx#`i t#Kv#bi Nv RwbZ cÖ`vn i#q#Q, t#v#l wWwRm A_@v AvBwjv#q#gi cÖ`vn i#q#Q, hK...#Zi cidvBwib wcv#K Amyweav i#q#Q Ges i#` RgvU evuavaq mgm`v i#q#Q Zv#`i mZK#Zvi ms#M t#b#b- t#me Kiv DwPz|

পানবিন্যাস :

thme tivMx#`i mwuq A_ev m#`n hy` t#ccwUK Avjmvi i#q#Q A_ev A#ö-Aw#Z`K i#` y#Y nq A_ev g#Svvi t`_#K g#v#ZK ai#Yi wKWbxi ywZ mwvaZ n#q#Q Zv#i t#b#b- e`envi Ki#eb bv| g#v#ZK fv#e hK...#Zi ywZMÖ` tivMxMY t#b#b- Øviv wPwKrmv Kiv#j Zv#`i hK...#Zi Kv#KvwiZv w#q#WZfv#e ch#teY Kiv#Z n#e|

অন্য ঔষধের সঙ্গে ব্যবহারে প্রতিক্রিয়া :

ছাই ইটেরিক্স : Ab`vb` bb-#U#ivBwvj cÖ`vn bvkK J#ai gZ t#b#b, WvBB#iwU-#i Kv#KvwiZv evavMÖ`i Ki#Z cv#i|

লিথিয়াম এবং ডিগক্সিন :

Ab`vb` bb-#U#ivBwvj cÖ`vn bvkK J#ai gZ t#b#b, wjw_qvg Ges wW#cvw-b GKm#½ e`envi Ki#j i#`i cövRgvq wjw_qvg Ges w#f#Mvw-b g#lv t#e#o th#Z cv#i|

অন্যান্য নন-স্টেরোইডাল প্রদাহ নাশক ঔষধ এবং স্টেরোইডস :

Ab`vb` bb-#U#ivBwvj cÖ`vn bvkK J#ai Ges t#U#ivBwM GK#i e`envi Ki#j cvk#©-cÖW-Zw#qv`i nvi t#e#o th#Z cv#i|

মাত্রাতিরিক্ত ব্যবহার :

t#b#b- g#lvwZwi#` e`env#ii d#j Bnvi t#k#Y tivaK#i M`vwU*K j`v#R Ges GKwU#U#UW P#K#öj Øviv wPwKrmv Kiv#Z n#e| RvUjZv Dck#gi Rb` jv#vYK cÖ`ZKv#K#i|

প্যাকেজের পরিমাণ :

v#Q 10 x 10 টি x 10 U`ve#iH|

কেমিস্ট ল্যাবরেটরীজ লিমিটেড

কলেজ রো, বরিশাল, বাংলাদেশ।



Composition:
Each film coated tablet contains Aceclofenac BP 100 mg

Description:
Painex contains aceclofenac that is a non-steroidal agent with anti-inflammatory and analgesic properties. Its mode of action is largely based on the inhibition of the enzyme cyclooxygenase that is involved in the production of prostaglandin. It also stimulates cartilage matrix (glycosaminoglycans) synthesis.

Indications :
Painex is indicated for the relief of pain and inflammation in osteoarthritis, rheumatoid arthritis and ankylosing spondylitis.

Dosage and Administration: By mouth
Adult : 1 tablet (100 mg) twice daily.

Children : There is no clinical data on the use of aceclofenac in children. So children should not be treated with painex.

Elderly: The pharmacokinetics of aceclofenac is not altered in elderly patients. Therefore it not considered necessary to modify the dose and dose frequency.

Hepatic insufficiency: The dose of aceclofenac should be reduced in patients with hepatic impairment. An initial daily dose of one tablet (100 mg) should be administered. **Renal insufficiency:** There is no evidence that the dose of painex needs to be modified in patients with mild renal impairment.

Side Effects:
Painex is generally well tolerated. The majority of the side effects are reversible and minor in nature. These include nausea, diarrhoea, dyspepsia, abdominal pain, pruritus, rash and occasionally dizziness.

Contraindication:
Painex is contraindicated in patients who are sensitive to aceclofenac or aspirin or other NSAIDs.

Use in Pregnancy:
There is no information on the safe use of aceclofenac during pregnancy. So Painex should not be administered during pregnancy unless there are compelling reasons for doing so. The lowest effective dose should be administered.

Use in Lactation:
There is no information about the secretion of aceclofenac in breast milk. So Painex should be avoided during lactation unless the potential benefits to the mother outweigh the possible risk to the children.

Precautions:
Painex should be administered with caution in patients with ulcerative colitis, crohn's disease, hepatic porphyria and coagulation disorders.

Warnings:
Painex should not be administered to patients with active or suspected peptic ulcer or gastrointestinal bleeding or moderate to severe renal impairment. Patients suffering from severe hepatic impairment must be monitored if they are treated with Painex.

Drug Interaction:
Diuretics: Painex like other NSAIDs may inhibit the activity of diuretics.
Quinolones: Convulsion may occur due to an interaction between quinolones and NSAIDs.

Lithium and digoxin: Painex like other NSAIDs may increase the plasma concentration of lithium and digoxin if taken together.

Other NSAIDs and steroids: Concomitant therapy of Painex with aspirin, other NSAIDs and steroids may increase the frequency of side effects.

Over Dosage:
In case of over dosage of Painex the absorption should be prevented as soon as possible by means of gastric lavage and treatment with activated charcoal. Supportive and symptomatic treatment should be given to relieve the complications.

Package Quantities:
10 x 10, 15 x 10 tablets in a box.

Manufactured by :
Chemist Laboratories Ltd.
College Row, Barishal, Bangladesh.